Generation of Flavour Dependent Lepton Asymmetries in the E$_6$SSM

S. F. King
University of Southampton, Southampton, SO17 1BJ, UK
R. Luo, D. J. Miller, R. Nevzorov
University of Glasgow, Glasgow, G12 8QQ, UK

We calculate flavour dependent lepton asymmetries within the E_6 inspired Supersymmetric Standard Model (E$_6$SSM). Our analysis reveals that the substantial lepton CP asymmetries can be induced even if $M_1 \simeq 10^6$ GeV.

arXiv:0810.0516v2 [hep-ph] 16 Oct 2008

1. INTRODUCTION

In models with heavy right–handed neutrinos lepton asymmetry can be dynamically generated via the out–of equilibrium decay of the lightest right–handed neutrino N_1. This asymmetry subsequently gets converted into the baryon asymmetry due to sphaleron interactions. In the standard model and its minimal supersymmetric extension the appropriate amount of baryon asymmetry can be induced only if the mass of the lightest right–handed neutrino $M_1 \gtrsim 10^9\,\mathrm{GeV}$. In this case reheat temperature has to be relatively high, i.e. $T_R > 10^9\,\mathrm{GeV}$, that results in an overproduction of gravitinos. To avoid gravitino problem T_R should be low enough $T_R \lesssim 10^{6-7}\,\mathrm{GeV}$. Here we study the generation of lepton CP asymmetries within the the Exceptional Supersymmetric Standard Model (E$_6$SSM) [1].

2. THE E$_6$SSM

The E$_6$SSM is based on the $SU(3)_C \times SU(2)_W \times U(1)_Y \times U(1)_N$ gauge group which is a subgroup of E_6. By definition $U(1)_N = \frac{1}{4}U(1)_\chi + \frac{\sqrt{15}}{4}U(1)_\psi$, where $E_6 \to SO(10) \times U(1)_\psi$, $SO(10) \to SU(5) \times U(1)_\chi$. To ensure anomaly cancellation the particle content of the E$_6$SSM is extended to include three complete 27 representations of E_6 [1]. The 27_i multiplets contain SM family of quarks and leptons, right–handed neutrino N^c_i, SM singlet field S_i which carry non–zero $U(1)_N$ charge, a pair of $SU(2)_W$–doublets H^d_i and H^u_i which have the quantum numbers of Higgs doublets and a pair of colour triplets of exotic quarks $\overline{D}_i$ and D_i which can be either diquarks (Model I) or leptoquarks (Model II). H^d_i and H^u_i form either Higgs or inert Higgs multiplets. In addition to the complete 27_i multiplets the low energy particle spectrum of the E$_6$SSM is supplemented by $SU(2)_W$ doublet L_4 and anti-doublet $\overline{L}_4$ from extra $27'$ and $\overline{27'}$ to preserve gauge coupling unification [1]. In the E$_6$SSM the right-handed neutrinos do not participate in the gauge interactions and therefore are expected to gain masses at some intermediate scale, while the remaining matter survives down to the electroweak scale. The part of the E$_6$SSM superpotential describing the interactions of N^c_i with other fields can be written as

$$W_N = h^N_{kxj}(H^u_k L_x)N^c_j + g^N_{kij} D_k d^c_i N^c_j\,, \tag{1}$$

where $x = 1, 2, 3, 4$ while $k, i, j = 1, 2, 3$. In the Model I $g^N_{kij} = 0$ whereas in the Model II all terms in Eq. (1) can be present.

3. CP ASYMMETRIES

The process of the lepton asymmetry generation is controlled by the flavour CP (decay) asymmetries. The CP asymmetries associated with the decays of N_1 and $\widetilde{N}_1$ into leptons and sleptons are given by

$$\varepsilon^k_{1,\,f} = \frac{\Gamma^k_{1f} - \Gamma^k_{1\bar{f}}}{\sum_{m,f'}\left(\Gamma^m_{1f'} + \Gamma^m_{1\bar{f}'}\right)}, \qquad \varepsilon^k_{\widetilde{1},\,f} = \frac{\Gamma^k_{\widetilde{1}f} - \Gamma^k_{\widetilde{1}\bar{f}}}{\sum_{m,\,f'}\left(\Gamma^m_{\widetilde{1}f'} + \Gamma^m_{\widetilde{1}\bar{f}'}\right)}, \tag{2}$$



Figure 1: Diagrams that give contribution to $\varepsilon^i_{1,\,\ell_k}$.

where f and f' correspond to either lepton or slepton field while Γ^k_{1f} and $\Gamma^k_{\tilde{1}f}$ are partial decay widths of $N_1 \to f + H^u_k$ and $\widetilde{N}_1 \to f + H^u_k$. At the tree level CP asymmetries vanish. The non–zero contributions to the CP asymmetries arise from the interference between the tree–level amplitudes of the lightest right–handed neutrino decays and one–loop corrections to them. Supersymmetry ensures that $\varepsilon^k_{1,\,\ell_k} = \varepsilon^k_{1,\,\widetilde{\ell}_k} = \varepsilon^k_{\widetilde{1},\,\ell_k} = \varepsilon^k_{\widetilde{1},\,\widetilde{\ell}_k}$. The tree–level and one–loop diagrams that contribute to the $\varepsilon^k_{1,\,\ell_k}$ are shown in Fig. 1. When $M_1 \ll M_2, M_3$ we get (see [2])

$$\varepsilon^k_{1,\,\ell_x} = -\frac{1}{8\pi A_1}\sum_{j=2,3}\mathrm{Im}\left[2A_j h^{N*}_{kx1}h^N_{kxj} + \sum_{m,\,y} h^{N*}_{my1}h^N_{mxj}h^N_{kyj}h^{N*}_{kx1}\right]\left(\frac{M_1}{M_j}\right), \tag{3}$$

where $A_j = \sum_{m,y} h^{N*}_{my1}h^N_{myj} + \frac{3}{2}\sum_{m,n} g^{N*}_{mn1}g^N_{mnj}$ if $j = 2,3$ while $A_1 = \sum_{m,y} h^{N*}_{my1}h^N_{my1}$.

The CP asymmetries associated with the decays of N_1 and $\widetilde{N}_1$ into exotic quarks (squarks) can be defined similarly to the lepton ones. The interference of the tree–level decay amplitude with the one–loop corrections (see Fig. 2) yields

$$\varepsilon^i_{1,\,D_k} = -\frac{1}{8\pi A_0}\sum_{j=2,3}\mathrm{Im}\left[2A_j g^N_{kij}g^{N*}_{ki1} + \sum_{m,\,n} g^{N*}_{mn1}g^N_{mij}g^N_{knj}g^{N*}_{ki1}\right]\left(\frac{M_1}{M_j}\right), \tag{4}$$

where $A_0 = \sum_{k,\,i} g^N_{ki1}g^{N*}_{ki1}$.

4. RESULTS AND CONCLUSIONS

To simplify our analysis we impose Z^H_2 symmetry under which all superfields except $H_d \equiv H^d_3$, $H_u \equiv H^u_3$ and $S \equiv S_3$ are odd. The Z^H_2 symmetry allows to avoid unacceptably large non-diagonal flavour transitions and reduces the part of the E$_6$SSM superpotential that describes the interactions of N^c_i with other superfields

$$W_N \to h^N_{3xj}(H_u L_x)N^c_j. \tag{5}$$

We also assume that the right–handed neutrino mass scale is relatively low ($M_1 \sim 10^6$ GeV), so that h^N_{3ij} are negligibly small ($i = 1,2,3$), and $M_3 \gg M_1, M_2$. In this approximation $\varepsilon^3_{1,\,L_4}$ is much larger than other CP asymmetries. In



Figure 2: Diagrams that give contribution to $\varepsilon^i_{1,\,D_k}$.

the considered case the maximal absolute value of $\varepsilon^3_{1,\,L_4}$ is given by

$$|\varepsilon^k_{1,\,\ell_4}| = \frac{3M_1}{8\pi M_2}|h^N_{342}|^2, \qquad (6)$$

where h^N_{342} is a coupling of the second lightest right–handed neutrino to H_u and L_4. From Eq. (6) one can see that in the considered case the substantial lepton decay asymmetry $\varepsilon^k_{1,\,\ell_4}$ ($\sim 10^{-6} - 10^{-4}$) can be induced even for $M_1 \simeq 10^6\,\mathrm{GeV}$ if $|h^N_{342}|$ vary from 0.01 to 0.1. At low energies the asymmetry generated in the L_4 sector gets converted into the ordinary lepton asymmetries via the decays of L_4. This suggests that in the E_6SSM successful thermal leptogenesis can be achieved without encountering the gravitino problem.

Acknowledgments

We would like to thank S. Antusch, C. D. Froggatt, L. B. Okun and D. Sutherland for fruitful discussions. RN acknowledges support from the SHEFC grant HR03020 SUPA 36878.

References

[1] S.F. King, S. Moretti, R. Nevzorov, "Theory and phenomenology of an exceptional supersymmetric standard model", Phys. Rev. D **73** (2006) 035009; S.F. King, S. Moretti, R. Nevzorov, "Exceptional supersymmetric standard model", Phys. Lett. B **634** (2006) 278; S.F. King, S. Moretti, R. Nevzorov, "Gauge Coupling Unification in the Exceptional Supersymmetric Standard Model," Phys. Lett. B **650** (2007) 57.

[2] S.F. King, R. Luo, D.J. Miller, R. Nevzorov, "Leptogenesis in the Exceptional Supersymmetric Standard Model: flavour dependent lepton asymmetries", arXiv:0806.0330 [hep-ph].